



AEROSOL OPTICAL AND CHEMICAL PROPERTIES WITHIN AND WITHOUT CLOUDS DURING AN AIRBORNE FIELD CAMPAIGN IN CENTRAL OKLAHOMA

Andrews, E., Lee, Y.-N., Alexander, M. L., Hubbe, J. M., and Ogren, J. A.

For presentation at the American Geophysical Union Fall Meeting San Francisco, CA December 10-14, 2007

[Eos Trans. AGU 88(52), Fall Meet. Suppl., Abstract A23D-1570]

Environmental Sciences Department/Atmospheric Sciences Division

Brookhaven National Laboratory

P.O. Box, Upton, NY www.bnl.gov

NOTICE: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

AEROSOL OPTICAL AND CHEMICAL PROPERTIES WITHIN AND WITHOUT CLOUDS DURING AN AIRBORNE FIELD CAMPAIGN IN CENTRAL OKLAHOMA

Andrews, E , betsy.andrews@noaa.gov , NOAA, NOAA/ESRL/GMD 325 Broadway, Boulder, CO 80305

Lee, Y , ynlee@bnl.gov , BNL, BNL PO Box 5000, Upton, NY 11973
Alexander, M L , lizabeth.alexander@pnl.gov , PNNL, PNNL PO Box 999, Richland, WA 99352
Hubbe, J M , john.hubbe@pnl.gov , PNNL, PNNL PO Box 999, Richland, WA 99352,
Ogren, J A , John.A.Ogren@noaa.gov , NOAA, NOAA/ESRL/GMD 325 Broadway, Boulder, CO 80305

The optical properties of aerosol particles are one of the controlling factors in determining direct aerosol radiative forcing. These optical properties depend on the chemical composition and size distribution of the aerosol particles, which can change due to various processes during the particles' lifetime in the atmosphere. Here we present preliminary results showing aerosol optical and chemical properties obtained during the CHAPS field campaign within cloud drops and outside of clouds. The Cumulis Humilis Aerosol Processing Study (CHAPS), sponsored by the DOE Atmospheric Science Program (ASP), took place in the vicinity of Oklahoma City in June, 2007. The intention of the study was to investigate the influence of clouds on aerosols and of aerosol on clouds. Duplicate sets of in-situ aerosol optical instruments were deployed on the ASP G-1 aircraft during the CHAPS campaign. One set of instruments was downstream of an isokinetic inlet designed to sample the ambient aerosol, the other set was downstream of a counterflow virtual impactor (CVI) designed to sample and dry cloud droplets so that the cloud drop nuclei could be studied. Each instrument set comprised a 3-wavelength particle soot absorption photometer (PSAP) and integrating nephelometer to provide spectral aerosol absorption, scattering and back-scattering and a particle counter to obtain aerosol number concentration. In addition, a time-of-flight aerosol mass spectrometer (ToF-AMS) was able to sample on either inlet to provide information about the non-refractory chemical composition of the aerosol and cloud drop residuals. The data presented here will describe both how aerosol optical properties change upstream and downstream of a mid-size conurbation (Oklahoma City) and how ambient aerosol optical properties differ from those of the cloud drop nuclei. These changes in aerosol optical properties will be placed in the context of differences in chemical composition derived from the ToF-AMS.